

Neocate® LCP

0+ months

Indications

For the dietary management of infants from birth (0+ months) with cows' milk allergy, multiple food protein allergy, eosinophilic oesophagitis and other indications where an amino acid-based formula is recommended.

Order information

Contact Nutricia Customer Care 1800 889 480

Flavour	Size	Code	Unit per carton
Unflavoured	400g	144542	6

Product Information

A nutritionally complete, amino acid based, powdered infant formula for special dietary use. Available on PBS.

Features

- Hypoallergenic.
- Nutritionally complete, suitable as a sole source of nutrition for infants under 1 year.
- Contains long chain polyunsaturated fatty acids (LCPs) (docosahexaenoic acid (DHA) & arachidonic acid (ARA)).
- Milk and soy protein free.

Important Notice

- Must be used under medical supervision after considering all feeding options, including breast feeding.
- For enteral use only, not for parenteral use.
- If using a tube feed, hang time should not exceed 4 hours.
- Suitable as a sole source of nutrition for infants under 1 year of age.
- Infants over 6 months should be offered food in addition to formula.

Direction of Use

- The quantity of feed should be determined by a clinician or dietitian only and is dependent on the age, body weight and medical condition of the patient.
- Each level scoop (4.5g) of Neocate® LCP requires 30ml of water to yield the recommended feed concentration (13.5% w/v).
- Follow the preparation instructions as stated on the can.

Storage

- Store unopened cans in a cool, dry place.
- Once opened, reseal can and store in a cool, dry place (not a refrigerator) and use within 4 weeks.

Feeding Guide

Approx. age of baby	Approx. weight of baby (kg)	Level scoops of Neocate LCP	Cooled boiled water (ml)	Feeds in 24 hours
Birth	3.5	3	90	6 – 7
2 weeks	4	4	120	6
2 months	5.5	5	150	5 – 6
3 months	6	6	180	5
5 months	7.5	7	210	4 – 5
7 months	8.5	7	210	4
10 months	9	8	240	4

Nutritional Information

Average contents	Per 100g	Per 100ml*
NUTRITION INFORMATION		
Energy	2062 kJ	278 kJ
	493 kcal	67 kcal
Protein Equivalent	13.3 g	1.8 g
	11 %E	
Carbohydrates	52.5 g	7.1 g
	42 %E	
Sugars	**4.6 g**	**0.62 g**
as Lactose	**0 g**	**0 g**
Total Fat	**25.6 g**	**3.5 g**
	47 %E	
Saturates	**10 g**	**1.3 g**
Polyunsaturates	**4.2 g**	**0.56 g**
of which LA	**3360 mg**	**453 mg**
of which ALA	**340 mg**	**45.6 mg**
of which DHA	**120 mg**	**16.6 mg**
of which AA	**120 mg**	**16.6 mg**
ω6:ω3 ratio	**7.58**	
Fibre	**0 g**	**0 g**
MINERALS		
Sodium	**217 mg**	**29.3 mg**
	9.4 mmol	**1.27 mmol**
Potassium	**540 mg**	**72.9 mg**
	13.8 mmol	**1.86 mmol**
Calcium	**571 mg**	**77.1 mg**
Phosphorus	**372 mg**	**50.2 mg**
Magnesium	**51.9 mg**	**7.01 mg**
Chloride	**392 mg**	**53.0 mg**
Ca:P ratio	**1.5:1**	
VITAMINS		
Vitamin A	431 µg-RE	58.2 µg-RE
Vitamin D	11.8 µg	1.6 µg
Vitamin E	11 mg	1.48 mg
Vitamin K	44.5 µg	6 µg
Vitamin C	52.7 mg	7.11 mg
Thiamin	0.59 mg	0.08 mg
Riboflavin	0.59 mg	0.08 mg

Niacin	4.09 mg	0.55 mg
Vitamin B_6	0.48 mg	0.06 mg
Folic acid	68.5 µg	9.24 µg
Vitamin B_{12}	1.38 µg	0.19 µg
Biotin	19.2 µg	2.6 µg
Pantothenic Acid	3.05 mg	0.41 mg
TRACE ELEMENTS		
Iron	7.4 mg	1.0 mg
Zinc	5.42 mg	0.73 mg
Manganese	49 µg	7 µg
Copper	420 µg	57 µg
Iodine	103 µg	14 µg
Molybdenum	11.3 µg	1.53 µg
Selenium	18.7 µg	2.53 µg
Chromium	10.8 µg	1.46 µg
OTHERS		
Choline	137 mg	18.5 mg
Inositol	110 mg	14.9 mg
L-carnitine	9.85 mg	1.33 mg
Taurine	32 mg	4.32 mg
Nucleotides	17.2 mg	2.33 mg
Osmolality	340 mOsmol/kgH_2O	

***At the recommended concentration of 13.5% w/v**
Abbreviations:
LA – Linoleic Acid
ALA – Alpha Linoleic Acid
DHA – Docosahexaenoic Acid
AA – Arachidonic Acid

Ingredients

Dried glucose syrup, refined vegetable Oils (high oleic sunflower oil, coconut oil, rapeseed oil, sunflower Oil), L-arginine L-aspartate, L-leucine, L-lysine acetate, L-glutamine, L-proline, emulsifier (E472c), L-valine, glycine, L-isoleucine, dipotassium hydrogen phosphate, L-threonine, L-phenylalanine, L-tyrosine, L-serine, L-histidine, L-alanine, calcium phosphate dibasic, tripotassiumcitrate, sodium chloride, calcium carbonate, choline bitartrate, L-cystine, calcium glycerophosphate, L-tryptophan, calcium citrate, magnesium chloride, oil from crypthecodinium cohnii, L-methionine, calcium phosphate tribasic, Oil from Mortierella alpina, magnesium L-aspartate, inositol, L-ascorbic acid, antioxidants (sunflower lecithin, ascorbylpalmitate), taurine, ferrous sulphate, zinc sulphate, L-carnitine, DL-alpha tocopherylacetate, uridine 5′-monophosphate sodium salt, cytidine-5'-monophosphate, nicotinamide, calcium D-pantothenate, inosine 5′-monophosphate sodium salt, adenosine-5'-monophosphate, guanosine 5′-monophosphate sodium salt, copper sulphate, thiamin mononitrate, riboflavin, pyridoxine hydrochloride, retinyl acetate, potassium iodide manganese sulphate, pteroylmonoglutamicacid, phytomenadione, sodium selenite, chromium chloride, sodium molybdate, D-biotin,

cholecalciferol, cyanocobalamin.

BREAST MILK IS BEST FOR BABIES: Professional advice should be followed before using an infant formula. Introducing partial bottle feeding could negatively affect breast feeding. Good maternal nutrition is important for breast feeding and reversing a decision not to breast feed may be difficult. Infant formula should be used as directed. Proper use of an infant formula is important to the health of the infant. Social and financial implications should be considered when selecting a method of feeding. Infant Formula product for special dietary use to be used under medical supervision. This product is not suitable for general use.FOR HEALTHCARE PROFESSIONALS ONLY.